

Donnish Journal of Entomology and Nematology
Vol 1(1) pp. 001-008 February, 2015.
<http://www.donnishjournals.org/djen>
Copyright © 2015 Donnish Journals

Original Research Article

The Incidence of the Protozoa *Malamoeba locustae* and *Gregarina granhami* in three Acridian Communities during Summer Month

*Abdel Rahman KM, El-Shazly MM and Ghazawy NA

Entomology Department, Faculty of Science, Cairo University, Giza, Egypt

Accepted 20th December, 2014.

The incidence of *Gregarina granhami* and *Malamoeba locustae* was investigated during summer 2012 among three wild grasshopper communities. Pooling the data of the three communities, it was found that about 38.5% of the grasshopper species hosted either *G. granhami* or *M. locustae*, and 15.28% of the individuals hosted one of them. The absence of such protozoa from *Chrotogonus lugubris* was attributed to an acquired immune response associated with feeding preference. It was assumed that these parasites are among exogenous density-dependent key factors regulating the grasshopper populations, and hence play a role in community stability in the wild acridid communities. The grasshopper communities were investigated at three study sites. The first site, Abu-Rauwâsh, is a countryside of Giza governorate. The second and third sites, Wadi El-Natroun and Kom Osheem, are located about 100 kilometers to the north and the south of the governorate; respectively. The first community consisted of 8 species; the most abundant was *Heteracris littoralis* (relative abundance 29.14%). The percentage of *G. granhami* in the community was 8.47% while *M. locustae* 27.45% of the population corresponding to 8.0% of the entire community. The second community consisted of 12 species; the most abundant was *H. littoralis* (relative abundance 22.49%). The percentage of *G. granhami* in the community was 6.89%, while *M. locustae* was recorded in 14.39% of *H. littoralis* population corresponding to 3.23% of the entire community. The third community consisted of 10 species; the most abundant was *Pyrgomorpha conica* (relative abundance 20.08%). The percentage of *G. granhami* in the community was 13.04%; it was recorded in *H. littoralis*, *P. conica* and *P. cognata*; again, *M. locustae* was recorded only in *H. littoralis* (28.47% of the population), corresponding to 5.66% of the entire community.

Keywords: Malamoeb, Gregarina, grasshopper, Orthoptera, Occupancy

INTRODUCTION

Parasitism, and other symbiotic activities are among the principal biological processes responsible for the transport and storage of material and energy, and the interactions of the organisms engaged in these activities may provide the pathways of feedback loops controlling the population regulation and community stability. Such extensive linkage between organisms has a profound impact on the organization in ecosystems through the food web (Price, 1999; El-Shazly, 2002).

El-Shazly and Shahpa (2006) analyzed the structure and host plant selection of an acridid community in Egypt and determined the diet breadth for each acridid species. In general, grasshopper community studies have recently been centered on the mechanisms driving community assembly (Rominger and Miller 2009), as well as on the analysis of the environmental factors controlling temporal and spatial

distribution of the investigated community (Rominger *et al.*, 2009, and Zhou *et al.*, 2012).

The principal pathogenic protozoa of locusts and grasshoppers are found in the groups amoeba, (Phylum: Sarcocystophora), gregarines (Phylum: Apicomplexa), and microsporidia (Phylum: Microspora); the most extensively studied group is the last one (Street and McGuire, 1990); primarily because of their potential as biological control agents (Abdel Rahman and Cagáñ 2001).

Concerning amoebic group, *M. locustae* is the only species of the genus *Malamoeba*; this species is characterized by a wide host range (Ernest and Baker 1982, Henry *et al.*, 1985), and it is responsible for amoebic diseases of grasshoppers (Street and McGuire, 1990). In contrast, some literatures suggest that host specificity is a general feature of gregarines; for instance, Clopton and Gold (1996) suggested that *Gregarina blattarum* comprised a complex of cryptic species marked by narrow host utilization rather than a single species persisting a broad array of cockroach taxa.

The amoeba *M. locustae* has been isolated more frequently from grasshoppers in laboratory cultures than from field collected grasshoppers. The isolation of this protozoan from laboratory cultures has been documented by King and Taylor (1936), Taylor and King (1937), Prinsloo (1960), Henry (1968) and Henry and Oma (1975). Among the above authors, Taylor and King 1937 described *M. locustae* from *Melanoplus differentialis*, *M. (mexicanus) sanguinipes* and *M. femurrubrum* raised in laboratory cultures in Iowa. Likewise, *G. garnhami* was described from an artificial population of *Schistocerca gregaria* raised in the Anti-Locust Research Center, London (Canning 1956).

The incidence of protozoan parasites on grasshopper communities is probably among the key factors controlling the community structure and species abundance. However, no attention has been paid, so far for the existence of these parasites in the wild grasshoppers communities; furthermore, Clopton and Gold (1996) pointed out that nearly 80% of the gregarine species have not been reported since their original description. Consequently, the aim of the present work is to investigate and compare the occurrence of *M. locustae* and *G. garnhami* in three acridid communities found in different habitats and separated by fairly large distances.

MATERIALS AND METHODS

Study area

Species were collected from non national park or other protected area, they live naturally on wild grasses close to the cultivated field owned by natives who allowed us to collect such grasshoppers. These specimens are not endangered or protected species. Locations of our study (GPS coordinates) are listed in the manuscript: Site I, Abu-Rauwāsh (30° 2' N; 31° 5' 42' E); Site II, Wadi El-Natroun (30° 16' N; 30° 02' E). Site III, Kom Osheem (29° 33' N 30° 53' E). For contact: Giza, El-Beheira and Fayom Governorates hotlines +20 2114, +20 45 3349848 and +20 84 6336577; respectively. These study sites (Fig. 1) were chosen to study the incidence of *M. locustae* and *G. garnhami* in grasshopper communities.

Site I, Abu-Rauwāsh district

A countryside of Giza governorate; it is an old land, cultivated with usual summer crops and irrigated with Nile water. The most abundant weed species in the sampling transect were *Cyperus rotundus* (L.) (Cyperaceae), *Cynodon 52 dactylon* (L.), *Polypogon monspiliensis* (L.), *P. monspiliensis* (L.) (Graminae), *Ammi majus* (L.) (Umbellifera), *Portulaca oleraceae* (L.) (Portulacaceae), *Alhagi maurorum* (Medic.) (Leguminosae), and *Brassica nigra* (L.) (Crucifera).

Site II, Wadi El-Natroun district

A depression in the western desert of Egypt located about 100 kilometers to the north of Giza governorate and belonging to El-Beheira governorate. Cultivated areas of this site where reclaimed land, irrigated with underground water. The abundant wild plants in the selected sampling transect were *C. rotundus* (L.) (Cyperaceae), *C. dactylon* (L.), *Avena fatue*, *Imperata cylindrical* (L.) (Graminae), *P. oleracea* (Portulacaceae), and *Eucalyptus* sp. (Myrtaceae).

Site III, Kom Osheem district

A reclaimed land cultivated mainly with olives and figs, and irrigated with Nile water; it is located about 100 kilometers to the south of Giza governorate and belonging to Fayom governorate. Beside the common weeds, *C. rotundus* (L.) (Cyperaceae), *C. dactylon* (L.), *Imperata cylindrical* (L.) (Graminae), other wild plants including *P. oleraceae* (L.) (Portulacaceae), *Alhagi maurorum* (Medic.) (Leguminosae), *Artiplex* sp. (Amaranthaceae) and *Schanginia baccata* (Chenopodiaceae) are common in this site.

Sampling

For each study site, one sampling transect about 2 meter width and 3000 meter length was designated. Care was taken to select the transects close to the borders of the cultivated fields where naturally growing plants are abundant. Because natural vegetation in Egypt is heterogeneous, patchy and fragmentary, care was taken to designate sampling transects with the most possible dense plant cover. The data were collected between June and August 2012.

The number of sampling points varied in the three sites according to the presence of low vegetation and grass patches; it ranged between 18 (in Wadi El-Natroun), and 27 sampling points (in Abu-Rauwāsh). The area of each sampling point was about 2x5 m along which grasshoppers were swept for 200 times using pest-net (net diameter 30 cm). The data of each sampling point is the average of 3 monthly samplings.

Examination

Collected insects were transferred to the laboratory. They were cut off at the fourth abdominal segment; then gut content, fat body and Malpighian tubes were crushed on a glass slide by a cover with a drop of distilled water and were examined by Leica microscope for seeking the microorganisms which were identified and photographed.

RESULTS AND DISCUSSION

Identification of the protozoa

The gregarine species in our study was considered as *Gregarina garnhami* (Fig. 2); while the amoebic one was considered as *Malamoeba locustae* (Fig.3) *G. garnhami* were identified according to Canning (1956) and Valigurova and Koudela (2008) based on the morphology of the free spores in the gut of the collected grasshoppers; on the other hand, *M. locustae* is the only species in the genus *Malameba* that is responsible for the amoebic disease of grasshoppers (Street and Mc Gure 1990). It was originally described by King and Taylor (1934) from Malpighian tubules of several *Melanoplus* species.

GRASSHOPPER COMMUNITIES

Site I (Abu-Rauwāsh):

A total of 8 grasshopper species and 1050 Individuals were recorded during the sampling period; the most abundant species was *H. littoralis* (relative abundance 29.14%); while the African migratory locust, *Locusta migratoria* (solitary phase) was the least abundant one (relative abundance 2.28%) (Table 1). Most populations exhibited peaks between July and August; except the native species, *A. aegyptium*, which was peaked in September (Fig.4). Both *M. locustae* and *G. garnhami* were recorded in *H. littoralis*, the former species was recorded in 27.25% of the collected specimens, corresponding to 8.0% of the community and the second in 4.28%.

However, the coexistence of the two parasites in the same individual was not observed; corresponding to 7.01 and 8.47% of the total community, respectively. *G. garnhami* was recorded in 10.21% of *A. aegyptium* population (1.81% of the community). Thus, we can state that 25% of the species, and 16.47% of the community were attacked by a protozoan species. Both *M. locustae* and *G. garnhami* were recorded in a wild population of *H. littoralis* for the first time. The former species, *G. garnhami*, was originally described from specimens of *S. gregaria*, *L. migratoria* and *A. aegyptim*; specimens of the latter species were sent to the author (Canning, 1956) from Egypt.

Table (1) The structure of acridian communities and the incidence of *Malamoeba locustae* (Mal.) and *G. garhami* (Gre)

Community (I) Abu-Rauwâsh						
Acridian species	Number collected	Relative abundance	No. infected		% infected	
			Mal.	Gre.	Mal.	Gre.
<i>Acrida pellucida</i>	127	12.09	-	-	-	-
<i>Acrotylus insubricus</i>	181	17.24	-	-	-	-
<i>Aiolopus thalassinus</i>	120	11.43	-	-	-	-
<i>Chrotogonus lugubris</i>	65	6.02	-	-	-	-
<i>Heteracris littoralis</i>	306	29.14	84	45	27.45	14.70
<i>Anacridium aegyptium</i>	186	17.71	-	19	-	10.21
<i>Locusta migratota</i>	24	2.28	-	-	-	-
<i>Truxalis nasuta</i>	41	3.90	-	-	-	-
∑ 8	∑ 1050		∑ 84	∑ 89		
Community (II) Wadi El-Natroun						
<i>Acrida pellucida</i>	61	5.06	-	-	-	-
<i>Acrotylus insubricus</i>	73	6.06	-	-	-	-
<i>Aiolopus thalassinus</i>	51	4.23	-	-	-	-
<i>Chrotogonus lugubris</i>	241	20.00	-	-	-	-
<i>Heteracris littoralis</i>	271	22.49	39	39	14.44	14.44
<i>Platypterna sp.</i>	25	2.07	-	-	-	-
<i>Pyrgomorpha conica</i>	240	19.92	-	10	-	4.17
<i>Pyrgomorpha cognata</i>	60	9.98	-	17	-	28.33
<i>Sphingonotus savignyi</i>	24	1.99	-	17	-	70.83
<i>Sphingonotus rubescens</i>	62	5.16	-	-	-	-
<i>Tropidopola longicornis</i>	51	4.23	-	-	-	-
<i>Truxalis nasuta</i>	46	3.82	-	-	-	-
∑ 12	∑ 1205		∑ 39	∑ 83		
Community (III) Kom Osheem						
<i>Acrida pellucida</i>	132	9.11	-	-	-	-
<i>Acrotylus insubricus</i>	270	18.63	-	-	-	-
<i>Aiolopus thalassinus</i>	177	12.21	-	-	-	-
<i>Chrotogonus lugubris</i>	62	4.28	-	-	-	-
<i>Heteracris littoralis</i>	288	19.87	82	38	28.47	13.19
<i>Platypterna</i>	52	3.59	-	-	-	-
<i>Pyrgomorpha conica</i>	291	20.08	-	96	-	32.99
<i>Pyrgomorpha cognata</i>	66	4.55	-	17	-	25.76
<i>Sphingonotus savignyi</i>	52	3.59	-	-	-	-
<i>Truxalis nasuta</i>	59	4.07	-	-	-	-
∑ 10	∑ 1449		∑ 82	∑ 189		

Table (2) The overall incidence of *Malamoeba locustae* (Mal.) and *Gregarina granhami* (Gre.) in grasshopper communities.

Acridian species	Collective percent infection in the three communities*					
	On the species level			On the communities level		
	Mal.	Gre.	Mal. + Gre.	Mal.	Gre.	Mal. + Gre.
<i>Acrida pellucid</i>	-	-	-	-	-	-
<i>Acrotylus insubricus</i>	-	-	-	-	-	-
<i>Aiolopus thalassinus</i>	-	-	-	-	-	-
<i>Anacridium aegyptium</i>	-	10.21	10.21	-	0.51	0.51
<i>Chrotogonus lugubris</i>	-	-	-	-	-	-
<i>Heteracris littoralis</i>	23.70	14.10	38.15	5.53	3.29	8.91
<i>Locusta migratota</i>	-	-	-	-	-	-
<i>Platypterna sp.</i>	-	-	-	-	-	-
<i>Pyrgomorpha conica</i>	-	19.96	19.96	-	2.86	2.86
<i>Pyrgomorpha cognate</i>	-	26.98	26.98	-	0.92	0.92
<i>Sphingonotus savignyi</i>	-	70.83	70.83	-	0.46	0.46
<i>Sphingonotus rubescens</i>	-	-	-	-	-	-
<i>Tropidopola longicornis</i>	-	-	-	-	-	-
<i>Truxalis nasuta</i>	-	-	-	-	-	-

*based on summation of data given in table (1)




Fig. 1: Map of Egypt showing the three different localities.



Fig. 2. *Gregarina garnhami*




Fig. 3. *Malamoeba locustae*




Fig. 4: Mean number of grasshoppers (No.) infected with *Malameoba locustae* (Mal.) and *Gregarina granhami* (Gre.) in Abu-Rauwâsh during summer months. Ac., *Acrida pellucida*; Ai., *Aiolopus thalassinus*; An., *Anacridium aegyptium*; Ar., *Acrotylus insubricus*; Ch., *Chrotogonus lugubris*; He., *Heteracris littoralis*; Lm., *Locusta migratoria*; Tn., *Truxalis nasuta*.




Fig. 5: Mean numbers of grasshoppers (No.) infected with *Malameoba locustae* (Mal.) and *Gregarina granhami* (Gre.) in Wadi El-Natroon during summer months. Ac., *Acrida pellucida*; Ai., *Aiolopus thalassinus*; Ar., *Acrotylus insubricus*; Ch., *Chrotogonus lugubris*; He., *Heteracris littoralis*; Pl., *Platypterna* sp.; Pg., *Pyrgomorpha conica*; Po., *Pyrgomorpha cognata*; Sr., *Sphingonotus rubescens*; Ss., *Sphingonotus savignyi*; Tr., *Tropidopola longicornis*; Tn., *Truxalis nasuta*.




Fig. 6: Mean number of grasshoppers (No.) infected with *Malameoba locustae* (Mal.) and *Gregarina granhami* (Gre.) in Kom Osheem during summer months. Ac., *Acrida pellucida*; Ai., *Aiolopus thalassinus*; Ar., *Acrotylus insubricus*; Ch., *Chrotogonus lugubris*; He., *Heteracris littoralis*; Pl., *Platypterna* sp.; Pg., *Pyrgomorpha conica*; Po., *Pyrgomorpha cognata*; Ss., *Sphingonotus savignyi*; Tr., *Tropidopola longicornis*.

SITE II: Wadi El-Natroun

The grasshopper community in sampling area of Wadi El-Natroun consisted of 12 species, four of them were parasitized with different levels of infections (Fig.5). This site seemed to be more rich in species than the other two sites, where 1205 specimens representing 12 species were collected (Table 1); this could be attributed to the loose nature of the sandy soil as well as the presence of heavy wild grass cover at the sampling site. Our field observations showed that many acridid species are connected to grasslands of every type from dry to relatively wet. The acridids that found in cultivated areas are usually connected to grass habitats along the borders of the fields. In a study on the ecology of *Acrida pellucida* in Egypt, Hafez and Ibrahim (1958) pointed out that adaptation of the genus *Acrida* to grasslands is an ancient one and that the evolution of the genus took place when grassland not only existed, but flourished, possibly during the Pliocene to the Pleistocene period.

A rapid glance at (Fig 5) is sufficient to show that the most abundant species was *H. littoralis* followed by *C. lugubris*, while the least abundant one was *S. savignyi*; the three species were represented by 22.49, 20 and 1.99% of the total number of specimens collected, respectively.

As in site I, *H. littoralis* was the only species infected by both *M. locustae* and *G. granhami*; three other species were infected by the gregarine parasite, namely *P. conica*, *P. cognata* and *S. savignyi*. The population of the last species, *S. savignyi*, had the heaviest infection where more than 70% of the sampled individuals hosted the gregarine parasite (Table1).

Site III: Kom Osheem

As shown in (Fig.6), the multispecies community in this site consisted of ten acridid species, with population peaked in midsummer (July-August); three species (*H. littoralis*, *P. conica* and *P. cognata*) hosted the protozoa. These species were represented by 1449 specimens. About 56.5% of the sample belonged to *P. conica*, *A. insubricus* and *H. littoralis* (Table 1). Poor representation of some species, e.g. *S. savignyi* in the collected sample could be attributed to innate or environmental factors such as reproductive capacity, seasonal history, phenology, availability of preferred food in the sampling area, suitability of soil for oviposition of a particular species etc. However, some species such as *H. littoralis* seemed to acquire a fair plasticity, having no definite preference for a particular habitat.

The overall incidence of the parasites on the three communities

The overall incidence of *M. locustae* (Mal.) and *G. granhami* (Gre.) in the three investigated grasshopper communities is given in (Table 2). Thirteen species were sampled from the three study sites; five species (about 38.5%) can host *G. granhami*, while *M. locustae* was found only in *H. littoralis*.

Gregarina free species could be resistant to the parasites. Thus *C. lugubris* is clean from such protozoa as it was collected from field several times since 1990 till now (personal observation); this could be attributed to an acquired immune response associated with feeding preference. In summer, this species feeds on garden purslane (*P. oleracea*: Portulacaceae) (Ibrahim, 1971) which contains flavonoids that has anti-protozoal activity (Upadhyay and Ahmad, 2012; Wang *et al.*, 2013). The susceptibility level of the host varied considerably among different acridid species, the more susceptible species was *S. savignyi*; more than 70% of the sampled individuals were infected with *G. granhami* (Table 2), and this may explain the low population level of the species in the communities (Table1).

Horizontal transmission via oral ingestion is the only reported route of amoebic disease infection in insects (Streett and Mc Guire, 1990). Among the conditions which contribute to the pathogenesis may be the absence of food and the tendency of grasshoppers and locusts to cannibalize upon dead or moribund individuals (Bidochka and Khachatourians, 1991). Indeed cannibalization of hanged individuals during molting has repeatedly been observed in grasshopper laboratory colonies (personal observations). This suggests that these parasites are among exogenous density-dependent key factors regulating the grasshopper populations, and hence play a role in the stability of wild acridid communities.

ACKNOWLEDGMENT

The authors wish to thank Dr. Hamed Fayed, professor of parasitology, zoology Department, Cairo University for his efforts in the identification of the parasites.

REFERENCES

- Abdel Rahman KM and Cagañ L (2001) Effect of *Nosema pyrausta* (Microsporidia, Nosematidae) on food consumption and utilization in the European corn borer, *Ostrinia nubilalis* (Lepidoptera, Crambidae). *Biologia* 56: 183-189.
- Ernst HP and Baker GL (1982) *Malameba locustae* (K & T) (Protozoa: Amoebidae) in field populations of Orthoptera in Australia. *J. Aust. Entomol. Soc.* 21: 295-296.
- Bidochka MJ and khachatourians GG (1991) Microbial and protozoan pathogens of grasshoppers and locusts as potential biocontrol agents. *Biocontrol Science and Technology* 1: 243-259
- Canning EU (1956) A New Eugregarine of Locusts, *Gregarina garhami* n.sp., parasitic in *Schistocerca gregaria* Forsk. *J. Protozool.* 3: 50-62.
- Clopton RE and Gold RE (1996) Host Specificity of *Gregarina blattarumvon* Siebold, 1839 (Apicomplexa: Eugregarinida) among Five Species of Domiciliary Cockroaches. *J. Inverteb. Pathol.* 67: 219-223.
- El-shazly MM (2002) Observations on oleander (*Nerium Oleander* L., Apocynaceae) ecosystem in Giza, Egypt. The 4th 173 International Conference on Urban Pests, Soth Carolina, USA: 225-233.
- El-shazly MM and Shahpa WME (2006) The structure and host plant selection of an acridid community on the edge between a desert- and an agro-ecosystem in Egypt; *Environmental Informatics Arch.* 4: 1-25.
- Hafez M and Ibrahim MM (1958) Ecological and biological studies on *Acrida pellucida* Klug, in Egypt. *Bull. Soc. Entomol. Egypte* 44: 451-476.
- Henry JE (1968) *Malameba locustae* and its antibiotic control in 179 grasshopper cultures. *J. Invertebr. Pathol.* 11: 224-233
- Henry JE and Oma EA (1975) Sulphonamide antibiotic control of *Malameba locustae* (King & Taylor) and its effect on grasshoppers. *Acrida* 4: 217-226
- Henry JE Wilson MC Oma EA and Fowler JL (1985) Pathogenic micro-organisms isolated from West African grasshoppers (Orthoptera: Acrididae). *Trop. Pest Manage.* 31:192-195.
- Ibrahim MM (1971) The ecology and biology of *Chrotogonus lugubris* Blanch. II. Food preference (Orthoptera: Pyrgomorphidae). *Bull. Soc. ent. Egypte* 52: 331-337.
- King RL and Taylor AB (1936) *Malpighamoeba locustae* n. sp. (Amoebidae), a protozoan parasite in the Malpighian tubes of grasshoppers. *Tran. Am. Microsc.* 55, 6-10.
- Price, PW (1999) The concept of the ecosystem. In: Huffaker, C. B., and Gutierrez, A. P., eds., *Ecological Entomology*, New York: John Wiley & Sons.
- Prinsloo HE (1960) Parasities Mikro-organisms by die Bruinsprinkaan *Locusta pardalina* (Walk). *S. Afr. Tydskr. Landbouwet*, 3: 551-560.
- Rominger AJ Miller TEX & Collins SL (2009) Relative contributions of neutral and niche-based processes to the structure of a desert grassland grasshopper community. *Oecologia* 161: 791-800.
- Street DA and McGuire MR (1990) Pathogenic diseases of grasshoppers. In Champman RF and Joern A. eds., *Biology of grasshoppers*. Wiley interscience. New York, pp: 483-516.
- Taylor AB and King RL (1937) Further studies on the parasitic amoebae found in grasshoppers. *Trans. Am. Microsc. Soc.* 56: 172-176.
- Valigurova A and Koudela B (2008) Morphological analysis of the cellular interactions between the eugregarine *Gregarina garhami* and the epithelium of its host, the desert locust *Schistocerca gregaria*. *Eur. J. Protistol.* 44: 197-207.
- Upadhyay RK and Ahmad S (2012) Ethno-medicinal plants and their pharmaceutical potential. *J. Pharm. Res.* 5: 2162-2173.
- Wang D Huang J Zhang Z Tian X Huang H. Yu Y Zhang G Ding J and Huang 205 R (2 0 1 3) Influences of *Portulaca oleracea* extracts on in vitro methane emissions and rumen fermentation of forage. *J. Food Agr. Environ.* 11: 4 8 3 - 4 8 8.
- Zhou W Wang K Zhao C and Zhang Q (2012) Analysis of spatial pattern among grasshopper and vegetation in Heihe based on GIS. *Physics Procedia* 33: 1261-1268.