

Original Research Article

Threshold Levels of Sperm Parameters Impacting on Pregnancy Rate in an Intrauterine Insemination Programme

Moez Kdous*, Kacem K, Ellouze S, Elloumi H, Khrouf M, Chaker A, Mnallah S and Zhioua A

Center of Reproductive Biology, Department of Gynecology and Obstetrics, Aziza Othmana Hospital, Tunis, University of Tunis El Manar, Tunisia.

Accepted 16th July, 2015.

Objectives: Intrauterine insemination (IUI) is frequently used as a first line strategy in the treatment of male and unexplained infertility. Threshold levels of sperm parameters associated with IUI success are controversial. The aim of this study was to evaluate the influence of sperm parameters on the outcome of IUI. **Study Design:** A prospective observational study. **Patients and methods:** This study included 295 IUI cycles. All IUI cycles were preceded by ovarian superovulation with clomiphene citrate 50 mg tablets orally twice daily for 5 days starting on the second day of menses and recombinant FSH 150 units IM on the 6th, 8th, and 10th day. Cycles were monitored by transvaginal ultrasound. The IUI was performed with a catheter 36 ± 4 hours after hCG injection. Sperm parameters before and after semen treatment for IUI were evaluated and correlated with pregnancy outcome. **Results:** A total of 29 clinical pregnancies were obtained, for a pregnancy rate per cycle of 9.83%. No pregnancy was obtained when less than one million spermatozoa were inseminated ($p = 0.022$). A statically significant increase in pregnancy rate was observed when normal sperm morphology was > 20% before semen treatment ($p = 0.01$) and > 25% after semen treatment ($p = 0.034$). NTSI (number of typical spermatozoa inseminated) and NTMSI (number of typical and mobile spermatozoa inseminated) significantly influence the clinical pregnancy rate. A 65% decrease in pregnancy rate was observed when the NTSI < 2 × 10⁶ compared to the NTSI ≥ 2 × 10⁶ (5.16% vs 15%, $p = 0.004$). Similarly, a 54% decrease in pregnancy rate was observed when the NTMS < 10⁶ compared to the NTMS ≥ 10⁶ (6.12% vs 13.5%, $p = 0.026$). **Conclusion:** IUI used for treating male factor infertility seems to have little chance of success when NMSI < 1 × 10⁶, NTSI < 2 million, NTMSI < 1 million. If these thresholds cannot be obtained, IVF should be recommended.

Keywords: Intrauterine insemination, Sperm parameters, Morphology, Motility, Pregnancy rate.

INTRODUCTION

Intrauterine insemination (IUI) is the simplest, least invasive and least expensive of the Assisted Reproductive Technologies (ART). Cervical infertility, male and unexplained infertility are still the major indications of this procedure [1]. However, the success rates of this technique are still low, suggesting that IUI should be retained only when it is medically indicated and biologically feasible.

Data from the literature are contradictory and do not allow to identify clearly which sperm parameters are associated with IUI success and which minimum threshold values should be accepted to use in daily practice in order to be able to establish a real guide of good practices for insemination [2]. The evaluation of our practices was therefore necessary so as to offer patients an adapted and personalized management. The

aim of our study was to determine the impact of the sperm parameters on the success of IUI.

PATIENTS AND METHODS

From January 2006 to December 2009, 295 couples (women younger than 40 years of age), all at their first attempt, completed 295 IUI cycles for male factor infertility in our department of biology and reproductive medicine. Institutional Review Board approval was obtained for this study. Before resorting to IUI, all women were subjected to testing for tubal patency by hysterosalpingography, and evaluating the pelvic anatomy by transvaginal ultrasonography. Laparoscopy was done when there was a possibility of pelvic adhesions or endometriosis in the HSG or transvaginal ultrasonography. The

assessment of the ovarian reserve was performed in all patients by a hormonal dosage done on day 3 of the cycle (FSH, LH, E2) associated to an ultrasound antral follicle count. Men had at least two semen analyses and microbiological tests before any treatment. All couples were tested for hepatitis B virus and hepatitis C virus before they were offered the IUI trial. Additional testing depended on any abnormalities observed.

Normal semen analyses were defined by the threshold values of the World Health Organization [3] (concentration $20 \times 10^6/\text{mL}$, total count 40×10^6 , progressive motility 50%, and typical morphology 30%). IUI was indicated only if at least 10^6 progressive motile spermatozoa were selected during a migration survival test. All semen samples were collected in the laboratory after 3 to 5 days of sexual abstinence. After liquefaction for 30 minutes at room temperature, volume, pH, sperm count, and progressive motility "a + b" were evaluated according to the WHO standard criteria.

Sperm concentration was performed with a hemocytometer on two separate preparations of the semen sample (dilution 1:20 in Ringer's solution). Sperm motility was determined by assessing at least five microscopic fields to classify at least 200 spermatozoa ($\times 4,000$ magnifications). The motility was graded progressive, non progressive, or immotile. Motile sperm were selected by a swim-up procedure. In all cases, the motile sperm fraction was washed twice by centrifugation, and the sperm pellet was suspended in 0.35 mL of Earle's balanced salt solution (Sigma-Aldrich Co., Ayrshire, United Kingdom) as a capacitating medium in all patients. Sperm were then counted, and progressive motility assessed. To analyze sperm morphology, smears were prepared from the whole ejaculated fraction and from the motile selected one.

The staining procedure was carried out according to the Sperm kit manufacturer's guidelines (Stain Enterprises, Onderstepoort, South Africa). The sperm smear was allowed to air-dry before being fixed in the formalin solution (fixative I) provided in the Sperm kit for 5 minutes at room temperature (23°C). Each slide was flooded with stain solution A for 2 minutes and then rinsed off with water. Each slide was then flooded with stain solution B and C each for 1 minute successively. The percentages of morphologically normal spermatozoa and of various sperm abnormalities were evaluated on 100 sperm at $\times 1,000$ magnification according to criteria of David and al. [4] modified by Jouannet and al. [5]

Ovarian stimulation was performed in all the patients with clomiphene citrate (Clomid®, Sanofi-Aventis, France) 100 mg / d from 2nd to 6th day of the cycle and recombinant FSH (GonalF®, Serono, France) 75 or 150 units on the 6th, 8th, and 10th day. Cycles were monitored by transvaginal ultrasound.

For the mean follicular volume and thickness of the endometrium on days 10, 12, and 14 of the cycle. Human chorionic gonadotropin 5,000 IU injection was given to induce ovulation when at least one follicle measured 18 mm or more. The IUI was performed with a catheter (Frydman catheter® (CCD, France) 36 ± 5 hours after hCG injection. All patients received 400 mg/day of dydrogesterone (Utrogestan®, International Besins, France) after insemination till the day of hCG testing. Serum hCG was determined 2 weeks after hCG injection for diagnosis of pregnancy. The outcome measure was the occurrence of clinical pregnancy, diagnosed by the visualization of fetal cardiac activity on ultrasound scan.

Data are expressed as the mean \pm SD or percentages. The χ^2 -test, Fisher's exact test, and Student's *t*-test were used for statistical analysis with SPSS software, version 15.0 for Windows (SPSS, Inc, Chicago, IL, USA). $P < .05$ was considered statistically significant.

RESULTS

The mean age of the women at IUI was 33.82 ± 5.074 years (range 23–40 years). According to the WHO criteria [3], various sperm abnormalities were encountered such as isolated oligospermia in 18 patients (6%), isolated asthenospermia in 12 patients (4%), isolated teratospermia in 65 patients (22%), oligoasthenospermia in 9 patients (3%), oligoteratospermia in 6 patients (2%), asthenoteratospermia in 36 patients (12%), and oligoasthenoteratospermia in 40 patients (13%).

Twenty-nine clinical pregnancies followed 295 IUI cycles, for a clinical pregnancy rate of 9.83 %. Spontaneous early abortions occurred in 11 pregnancies (3.7%), and twin pregnancies occurred in 4 patients (1.3%).

Table 1 shows no significant difference in: total sperm concentration, motility 'a', 'b' and "a + b" before and after sperm treatment, between patients who achieved pregnancy and those who failed. Only normal sperm morphology was significantly associated with pregnancy rate ($p=0.001$ before sperm treatment and 0.005 after)

Table 2 shows a statically significant increase in pregnancy rate when normal sperm morphology was $> 20\%$ before semen treatment ($p = 0.01$) and $> 25\%$ after semen treatment ($p = 0.034$).

Comparison of NMSI (number of motile spermatozoa inseminated), NTSI (number of typical spermatozoa inseminated) and NTMSI (number of typical and motile spermatozoa inseminated) between pregnant and non-pregnant groups showed no significant differences (table 3).

No pregnancy was obtained when less than one million spermatozoa were inseminated ($p = 0.022$). Above one million, there is no statistical difference in pregnancy rates between subgroups $1 \leq \text{NMSI} < 5$; $5 \leq \text{NMSI} < 10$ and $\text{NMSI} \geq 10$. Beyond 10 million, our results showed a slight decline (though not significant).

NTSI and NTMS significantly influence the occurrence of clinical pregnancy. A 65% decrease in pregnancy rate was observed when the $\text{NTSI} < 2 \times 10^6$ compared to the $\text{NTSI} \geq 2 \times 10^6$ (5.16% vs 15%, $p = 0.004$). Similarly, a 54% decrease in pregnancy rate was observed when the $\text{NTMS} < 10^{6\text{m}}$ compared to the $\text{NTMS} \geq 10^6$ (6.12% vs 13.5%, $p = 0.026$). (Table 4).

DISCUSSION

There is good evidence in literature that intrauterine insemination (IUI) is the best first line treatment and most cost-effective procedure for moderate male factor subfertility. It seems very difficult to identify individual semen parameters predicting the likelihood of pregnancy after IUI. This can be explained by a lack of standardization of semen analysis, but many other methodological variables may also influence IUI success rates, such as the patient selection, type of ovarian stimulation and number of inseminations per cycle [6].

The majority of studies attempting to find which sperm characteristics correlated better with cycle outcome included multiple female infertility problems as well. Hence, the results might be biased by the other infertility etiologies that were treated in parallel. In this study, only the male factor problems were recruited to make the differences in sperm parameters more significant. Twenty-nine clinical pregnancies were achieved after 295 IUIs (9.83%). In a review of 17 papers about IUI with ovarian stimulation by hMG, Wainer et al. [7] reported 274 pregnancies in 2,223 IUI cycles, for a pregnancy rate per cycle of 12.33%.

Table 1. Sperm parameters values observed in both pregnant and non-pregnant groups before and after semen treatment

	Before semen treatment			After semen treatment		
	Non pregnant (n=266)	Pregnant (n=29)	P	Non pregnant (n=266)	Pregnant (n=29)	P
Concentration ($10^6/ml$)	68.16±18.1	53.45±17.8	.288	24.26±10.5	25.59±11.2	.732
Motility « a » (%)	13.44±6.3	16.21±7.2	.132	36.02±8.3	41.38±9.1	.111
Motility « b » (%)	29.52±15.4	30.52±14.6	.592	35.17±10.3	34.66±10.5	.806
Motility « a+b » (%)	42.96±20.6	46.72±19.8	.178	73.12±18.6	76.03±19.1	.121
Typical form (%)	17.98±7.4	24.93±8.2	.001	22.15±8.6	29.76±7.7	.005

Table 2. Pregnancy rate correlated to sperm parameters values before and after semen treatment

	Before semen treatment				After semen treatment			
	%	cycles	Pregnancy n (%)	P	%	cycles	Pregnancy n (%)	P
Motility « a » (%)	<15	149	10 (6.7)	.052	<35	128	9 (7.03)	.111
	≥15	146	19 (13.01)		≥35	167	20 (11.97)	
Motility « a+b » (%)	<50	170	15 (8.82)	.314	<75	129	12 (9.3)	.474
	≥50	125	14 (11.2)		≥75	166	17 (10.2)	
Typical form (%)	<20	167	10 (5.9)	.010	<25	154	10 (6.4)	.034
	≥20	128	19 (14.84)		≥25	141	19 (13.4)	

Table 3. Comparison of NMSI, NTSI and NTMSI between pregnant and non-pregnant groups

	Non pregnant (n=266)	Pregnant (n=29)	P
NMSI (10^6)	5.3	5.45	.877
NTSI (10^6)	3.42	3.45	.977
NTMSI (10^6)	1.65	1.34	.471

NMSI: number of motile spermatozoa inseminated; **NTSI:** number of typical spermatozoa inseminated; **NTMSI:** number of typical and motile spermatozoa inseminated

Table 4. Pregnancy rate correlated to NMSI, NTSI, NMTSI

	cycles	Pregnancy n (%)	P
NMSI (10^6)	<1	35	0 (0)
	1≤N<5	119	14 (11.76)
	5≤N<10	99	11 (11.11)
	N≥10	42	4 (9.52)
NTSI (10^6)	<2	155	8 (5.16)
	≥2	140	21 (15)
NTMSI (10^6)	<1	147	10 (6.12)
	≥1	148	19 (13.5)

NMSI: number of motile spermatozoa inseminated, **NTSI:** number of typical spermatozoa inseminated;

NTMSI: number of typical and motile spermatozoa inseminated;

^a a 65% decrease in pregnancy rate; ^b a 54% decrease in pregnancy rate

It has previously been demonstrated that, the number of motile spermatozoa inseminated (NMSI) and sperm morphology are the most valuable sperm parameters to Predict IUI outcome. The minimum recommended number of motile sperm after semen treatment in various reports varies from 0.8 to 10×10^6 [8, 9, 10 and 11]. In the study of Badaway and al [12] the pregnancy rate per cycle was significantly lower in the subgroups with number of motile sperm $<5 \times 10^6$ (5.55% vs. 24.8%). This was in agreement with Francavilla and al. [13], who found a significant increase in success rate when the total motile sperm count exceeded 5×10^6 .

Van Weert and al [14] in a meta-analysis of 16 studies reported a cut-off of NMSI between 0.8 and 5 million. In this study, the number of motile sperm inseminated (NMSI) was not significantly different in the cycles resulting in pregnancy compared to the failed cycles. But, when (NMSI) was $<1 \times 10^6$,

no pregnancy was obtained in our study ($p < 0.05$). However, the low number of attempts with less than 1 million NMSI generated a sampling bias limiting the relevance of this parameter. Beyond 1 million, the NMSI does not seem to influence pregnancy rate significantly. But, when NMSI exceeded 10 million, our results showed a slight non-significant decrease in pregnancy rate, which could be related to excessive production of free radicals by the spermatozoa. Most authors recommend IVF when the number of motile sperm is $<1 \times 10^6$ [10, 11, 15 and 16].

Some authors suggested using the total number of motile spermatozoa of the initial sperm count (before semen treatment) as the criterion for choosing between IUI and IVF, and have recommended threshold values ranging from 5 to 10×10^6 [17, 18]. In other studies, total sperm motility before semen treatment was mentioned with a cut-off level between

30 and 50% [19, 20, 21 and 22]. The usefulness of this criterion, however, is limited by the variability of quality from one ejaculation to another, as well as the variable results of sperm preparations. Our study shows no significant difference in sperm motility before sperm treatment, between patients who achieved pregnancy and those who failed

Sperm morphology using strict criteria is well known as one of the best predictors of IVF outcome [23, 24 and 25]. Sperm morphology has also turned out to be a good predictive parameter in IUI. In a meta-analysis, a significant improvement in pregnancy rates above the 4% threshold for strict criteria was described [14]. Universally, the most common accepted classification system used for sperm morphology is the WHO criteria [19]. The threshold of spermatozoa with normal morphology below which IVF is recommended varies widely in the literature from 4% to 50% [27, 28 and 29]. The present analysis supported these results (thresholds of 20% in the initial sperm and 25% after semen treatment)

However, the implication of sperm morphology alone for choice and outcome of IUI is controversial. Many retrospective and prospective studies showed that sperm morphology alone, before or after preparation, did not help to predict IUI results [30, 31 and 32]. The evaluation of number of motile sperm in conjunction with sperm morphology might represent an important deliberation for clinical decision making. According to Wainer and al [7] For a normal morphology sperm rate <30%, the clinical pregnancy rate was 4.62% when the number of motile sperm inseminated (NMSI) was $<5 \times 10^6$ and 9.45% when the NMSI was $>5 \times 10^6$, and this difference was statistically significant, suggesting a partial compensation of quality by the amount of spermatozoa. The study of Badawy and al [12] showed that above 5 million motile spermatozoa inseminated, the clinical pregnancy rate per cycle was 5.3% when the normal morphology sperm rate (NTSI) rate <30% and 18.42% when $>30\%$.

Our study supported these findings and showed that the NTSI and the NTMSI significantly influence the pregnancy rate in IUI. Thus, when the NTSI was less than 2 million, IUI proved to be an inefficient ART indication with a pregnancy rate of only 5.16%. In contrast, when the NTSI was greater than or equal to 2 million pregnancy rate increases to 15% ($p = 0.04$). Similarly, when the NTMSI was less than 1 million, the success rate of IUI was 6.12%. This rate increased to 13.5% when the NTMSI was greater than or equal to 1 million ($p = 0.026$).

CONCLUSION

In view of these results, IUI used for treating male factor infertility seems to have little chance of success when NMSI $<1 \times 10^6$, NTSI <2 million, NTMSI <1 million. If these thresholds cannot be obtained, IVF should be recommended.

REFERENCES

- Marchetti C and Dewailly D. Intrauterine insemination: indications and methods. *Rev Prat* 2006; 56(5): 500-6.
- Royère D. Insémination intra-utérine : état des lieux chez l'humain. *Gynécol Obstet Fertil* 2004 ; 32(10) 873-9.
- World Health Organization. Laboratory manual for the examination and processing of human semen. Fourth edition. 1999
- G. David, J.P. Bisson, F. Czyglik, P. Jouannet, C. Gernigon. Morphological abnormalities of the human sperm. Proposition for a classification system. *J Gynecol Obstet Biol Reprod* 1975 ; 4 : 17-36
- Jouannet P, Ducot B, Feneux D, Spira A. Male factors and the likelihood of pregnancy in infertile couples. Study and sperm characteristics. *Int J Androl* 1988; 11: 379-384
- Ombelet W, Deblaere K, Bosmans E, and al. Semen quality and intrauterine insemination. *Reprod Bio Med Online* 2003 4 : 485-492.
- Wainer R, Albert M, Dorion A, Bailly M, Bergère M, Lombroso R, and al. Influence of the number of motile spermatozoa inseminated and of their morphology on the success of intrauterine insemination. *Hum Reprod* 2004; 19: 2060-2065
- Dodson WC, Haney AF. Controlled ovarian hyperstimulation and intrauterine insemination for treatment of infertility. *Fertil Steril* 1991; 55: 457-467
- Berg U, Brucker C, Berg FD. Effect of motile sperm count after swim-up on outcome of intrauterine insemination. *Fertil Steril* 1997; 67: 747-750
- Van Voorhis BJ, Barnett M, Sparks AET, Syrop CH, Rosenthal G, Dawson J. Effect of the total motile sperm count on the efficacy and cost-effectiveness of intrauterine insemination and in vitro fertilization. *Fertil Steril* 2001; 75: 661-668
- Miller DC, Hollenbeck BK, Smith GD, Randolph JF, Christman GM, Smith YR, and al. Processed total motile sperm count correlates with pregnancy outcome after intrauterine insemination. *Urology* 2002: 497-501
- Badawy, Elnashar A, Eltotongy M. Effect of sperm morphology and number on success of intrauterine insemination. *Fertil Steril* 2009; 91:777-781.
- Francavilla F, Romano R, Santucci R, Poccia G. Effect of sperm morphology and motile sperm count on outcome of intrauterine insemination in oligozoospermia and/or asthenospermia. *Fertil Steril* 1990, 53: 892-897
- Van Waart J, Kruger TF, Lombard CJ, Ombelet W. Predictive value of normal sperm morphology in intra uterine insemination (IUI): a structured literature review. *Hum Reprod Update* 2001; 7(5): 495-500.
- Brasch JG, Rawlins R, Tarchala S, Radwanska E. The relations between total motile sperm count and the success of intrauterine insemination. *Fertil Steril* 1994; 62: 150-154
- Horvath P, Bohrer M, Shelden R, Kemmann E. The relationship of sperm parameters to cycle fecundity in superovulated woman undergoing intrauterine insemination. *Fertil Steril* 1989, 52: 288-294
- Campana A, Sakkas D, Stalberg A, Grace Bianchi P, Comte I, Pache T, and al. Intrauterine insemination: evaluation of the results according to the woman's age, sperm quality, total sperm count per insemination and life table analysis. *Hum Reprod* 1996; 11: 732-736
- Elvan Koyun Ok, Ömer Erbil Doğan, Recep Emre Okay, and Bülent Gülekli. The effect of post-wash total progressive motile sperm count and semen volume on pregnancy outcomes in intrauterine insemination cycles: a retrospective study. *J Turk Ger Gynecol Assoc* 2013; 14(3): 142-145.
- Shulman A, Hauser R, Lipitz S and al. Sperm motility is a major determinant of pregnancy outcome following intrauterine insemination *Journal of Assisted Reproduction and Genetics* 1998; 15: 381-385.
- Dickey RP, Pyrzak R, Lu PY and al. Comparison of the sperm quality necessary for successful intrauterine insemination with World Health Organization threshold values for normal sperm. *Fertil Steril* 1999; 71, 684-689.
- Montanaro GM, Kruger TF, Coetzee K and al. Stepwise regression analysis to study male and female factors impacting on pregnancy rate in an intrauterine insemination programme. *Andrologia* 2001 ; 33 : 135-141.
- Lee RK-K, Hou J-W, Ho H-Y and al. Sperm morphology analysis using strict criteria as a prognostic factor in intrauterine insemination. *International Journal of Andrology* 2002; 25: 277-280.
- S. Cao S, C. Zhao C, J. Zhang J, X. Wu X, L. Zhou L, X. Guo X and al. A minimum number of motile spermatozoa are required for successful fertilisation through artificial intrauterine insemination with husband's spermatozoa. *Andrologia* 2014; 46 (5): 529-534.
- Saucedo de la Llata E., Moraga Sanchez M. R, Batiza Resendiz V, Galache Vega P, Santos Haliscak R, Hernandez Ayup S. Influence of sperm morphology on results of intrauterine insemination. *Ginecol Obstet Mex* 2003; 71: 455-9

- [25] Sun Y, Li B, Fan LQ et al. Does sperm morphology affect the outcome of intrauterine insemination in patients with normal sperm concentration and motility? *Andrologia* 2012; 44(5): 299–304
- [26] Kruger TF, Menkveld R, Stander FSH, Lombard CJ, van Zyl J, Smith K. Sperm morphologic features as a prognostic factor in in vitro fertilization. *Fertil Steril* 1986; 46: 1118–1123
- [27] Sakhel K, Abozaid T, Schwarck S, Ashraf M, and al Semen parameters as determinants of success in 1662 cycles of intrauterine insemination Effective controlled ovarian hyperstimulation *Fertil Steril* 2005; 84:248-249
- [28] Haim D, Leniaud L, Porcher R, Martin-Pont B, Wolf JP, Sifer C. Prospective evaluation of the impact of sperm characteristics on the outcome of intrauterine inseminations. *Gynecol Obstet Fertil* 2009; 37:229-235.
- [29] Berker B, Sükür YE, Kahraman K Atabekoglu CS, Sonmezer M, Ozmen and al Absence of Rapid Linear and Progressive Motile Spermatozoa "Grade A" in Semen Specimens: Does It Change Intrauterine Insemination Outcomes? *Urology* 2012; 80(6): 1262-6
- [30] Hassa H, Aydin Y, Oge T, Ozatik O, Taplamacioglu F. Predictive factors for clinical pregnancy in 354 unexplained infertile couple in their first intrauterine insemination cycle. *Fertil Steril* 2012; 98: 160
- [31] Matorras R, Corcostegui B, Perez C, Mandiola M, Mendoza R, Rodriguez-Escudero FJ. Sperm morphology analysis (strict criteria) in male infertility is not a prognostic factor in intrauterine insemination with husband's sperm. *Fertil Steril* 1995, 63: 608–611
- [31] Karabinus DS, Gelety DS. The impact of sperm morphology evaluated by strict criteria on intrauterine insemination success. *Fertil Steril* 1997; 67: 536–541